VOL. 666, NOS. 1 + 2 JOURNAL OF CHROMATOGRAPHY A APRIL 22, 1994

CONTENTS

CHIRAL SEPARATIONS —FUNDAMENTAL ASPECTS AND APPLICATIONS

Preface
by W. Lindner (Graz, Austria) 1

$\pi-\pi$ Type chiral selectors; Pirkle type CSPs

Evolution of chiral stationary phase design in the Pirkle laboratories (Review)
by C.J. Welch (Morton Grove, IL, USA) 3

Evaluation of Pirkle-type chiral stationary phases by liquid and supercritical fluid chromatography. Influence of the spacer length and the steric hindrance in the vicinity of the stereogenic centre
by N. Bargmann-Leyder (Paris, France), J.-C. Truffert and A. Tambuté (Vert-le-Petit, France) and M. Caude (Paris, France) 27

Diphenylethanediamine derivatives as chiral selectors. III. Comparison of four new diastereomeric chiral stationary phases prepared by addition of mono-3,5-dinitrobenzoyldiphenylethanediamine derivatives to optically pure epoxy silica
by G. Uray, N.M. Maier and W. Lindner (Graz, Austria) 41

Chiral selectors with chelating properties

Chiral selectors with chelating properties in liquid chromatography: fundamental reflections and selective review of recent developments (Review)
by V.A. Davankov (Moscow, Russian Federation) 55

Enantiomeric separation of dansyl- and dabsylamino acids by ligand-exchange chromatography with (*S*)- and (*R*)-phenylalaninamide-modified silica gel
by B. Galli, F. Gasparrini, D. Misiti and C. Villani (Rome, Italy) and R. Corradini, A. Dossena and R. Marchelli (Parma, Italy) 77

New chemically bonded chiral ligand-exchange chromatographic stationary phases
by G. Gübitz, S. Mihellyes, G. Kobinger and A. Wutte (Graz, Austria) 91

Unusual peak distortion in ligand-exchange chromatography of enantiomers under overloaded conditions
by A. Kurganov (Moscow, Russian Federation and Mainz, Germany), V. Davankov (Moscow, Russian Federation), K. Unger (Mainz, Germany) and F. Eisenbeiss and J. Kinkel (Darmstadt, Germany) 99

Gas chromatographic separation of enantiomers

Enantiomer separation by gas chromatography on chiral stationary phases (Review)
by V. Schurig (Tübingen, Germany) 111

Enantioselective separations by capillary gas chromatography on derivatized cyclodextrins. III. Separation of some racemic 2,2-dialkyl-4-alkoxycarbonyl-1,3-dioxolane derivatives on 2,3-di-O-acetyl-6-O-*tert.*-butyldimethylsilyl-β- and -γ-cyclodextrins
by K. Jaques, W.M. Buda, A. Venema and P. Sandra (Ghent, Belgium) 131

Cyclodextrin derivatives in the gas chromatographic separation of racemic mixtures of volatile compounds. VII. The use of 2,6-di-O-methyl-3-O-pentyl-β-cyclodextrin diluted in phases with different polarity in the separation of racemates in complex mixtures
by C. Bicchi, A. D'Amato and V. Manzin (Turin, Italy) and A. Galli and M. Galli (Legnano, Milan, Italy) . . . 137

Examination of the enantioselectivity of wall-immobilized cyclodextrin copolymers in capillary gas chromatography
by Y. Tang, Y. Zhou and D.W. Armstrong (Rolla, MO, USA) 147

Chiral compounds of essential oils. XV. Stereodifferentiation of characteristic compounds of *Mentha* species by multi-dimensional gas chromatography
by B. Faber, A. Dietrich and A. Mosandl (Frankfurt, Germany) 161

Proteins and peptides as chiral selectors

Proteins and peptides as chiral selectors in liquid chromatography (Review)
by S.G. Allenmark (Göteborg, Sweden) and S. Andersson (Linköping, Sweden) 167

Dynamic modification of the chiral bonding properties of a CHIRAL-AGP column by organic and inorganic additives. Separation of enantiomers of anti-inflammatory drugs
by J. Hermansson and I. Hermansson (Hägersten, Sweden) 181

Optical resolution by electrokinetic chromatography using ovomucoid as a pseudo-stationary phase
by Y. Ishihama, Y. Oda, N. Asakawa, Y. Yoshida and T. Sato (Ibaraki, Japan) 193

Separation of enantiomers on a lysozyme-bonded silica column
by J. Haginaka, T. Murashima and C. Seyama (Nishinomiya, Japan) 203

Changes in chiral selectivity with temperature for an ovomucoid protein-based column
by K.M. Kirkland and D.A. McCombs (Wilmington, DE, USA) 211

Stereo-recognition in life sciences, impact of chromatographic methods, application in pharmacology, etc.

Enantioselective high-performance liquid affinity chromatography as a probe of ligand–biopolymer interactions: an overview of a different use for high-performance liquid chromatographic chiral stationary phases
by I.W. Wainer (Montreal, Canada) 221

Investigations on the *in vitro* racemization of thalidomide by high-performance liquid chromatography
by B. Knoche and G. Blaschke (Münster, Germany) 235

Enantioselective assay for the determination of nisoldipine in dog, rat and mouse plasma by chiral microbore high-performance liquid chromatography combined with gas chromatography–mass spectrometry
by D. Zimmer and V. Muschalek (Wuppertal, Germany) 241

Enantiomer resolution of camazepam and its derivatives and enantioselective metabolism of camazepam by human liver microsomes
by X.-L. Lu and S.K. Yang (Bethesda, MD, USA) 249

Liquid chromatographic determination of D- and L-amino acids by derivatization with *o*-phthaldialdehyde and chiral thiols. Applications with reference to biosciences
by H. Brückner, S. Haasmann, M. Langer, T. Westhauser and R. Wittner (Stuttgart, Germany) and H. Godel (Waldbronn, Germany) 259

Stereoselective high-performance liquid chromatographic assay of (±)-delmopinol in plasma using solid-phase extraction, a chiral derivatizing agent and electrochemical detection
by G. Egginger, E. Blaschke and W. Lindner (Graz, Austria) and A.-M. Olsson (Malmö, Sweden) 275

Enantioselective high-performance liquid chromatographic method for the determination of methadone in serum using an AGP and a CN column as chiral and analytical column, respectively
by K. Kristensen and H.R. Angelo (Copenhagen, Denmark) and T. Blemmer (Herlev, Denmark) 283

Chromatographic resolution and pharmacological investigation of ICI 118551, a new β_2-blocker
by M.G. Quaglia, N. Desideri, E. Bossù, A. Farina and L.A. Morrone (Rome, Italy) 289

Separation of enantiomers by electrophoretic techniques

Separation of enantiomers by capillary electrophoretic techniques (Review)
by S. Terabe (Hyogo, Japan), K. Otsuka and H. Nishi (Osaka, Japan) 295

Packed-capillary electrochromatographic separation of the enantiomers of neutral and anionic compounds using β-cyclodextrin as a chiral selector. Effect of operating parameters and comparison with free-solution capillary electrophoresis
by S. Li and D.K. Lloyd (Montreal, Canada) 321

Capillary zone electrophoresis for separation of drug enantiomers using cyclodextrins as chiral selectors. Influence of experimental parameters on separation
by S. Pálmarsdóttir and L.-E. Edholm (Lund, Sweden) 337

Chiral separations of basic and acidic compounds in modified capillaries using cyclodextrin-modified capillary zone electrophoresis
by D. Belder and G. Schomburg (Mülheim an der Ruhr, Germany) 351

Enantiomeric separations in capillary zone electrophoresis using a chiral crown ether
by R. Kuhn (Reutlingen, Germany) and C. Steinmetz, T. Bereuter, P. Haas and F. Erni (Basle, Switzerland) . . 367

Chiral separation by capillary zone electrophoresis of an optically active drug and amino acids by host–guest complexation with cyclodextrins
by A. Werner, T. Nassauer, P. Kiechle and F. Erni (Basle, Switzerland) 375

Separation of enantiomers by supercritical fluid chromatography

Chiral separations performed by supercritical fluid chromatography (Review)
by P. Petersson and K.E. Markides (Uppsala, Sweden) 381

Chiral separations by packed-column super- and subcritical fluid chromatography
by K. Anton, J. Eppinger, L. Frederiksen and E. Francotte (Basle, Switzerland) and T.A. Berger and W.H. Wilson (Wilmington, DE, USA) 395

Hydrocarbon derivatives, amides and amines, as chiral selectors

Resolution by high-performance liquid chromatography using polysaccharide carbamates and benzoates as chiral stationary phases (Review)
by Y. Okamoto and Y. Kaida (Nagoya, Japan) 403

Enantiomeric separation of chiral sulphoxides. Screening of cellulose-based sorbents with particular reference to cellulose tribenzoate
by E. Küsters, V. Loux, E. Schmid and Ph. Floersheim (Basel, Switzerland) 421

Influence of temperature on chiral high-performance liquid chromatographic separations of oxazepam and Prominal on chemically bonded β-cyclodextrin as stationary phase
by K. Cabrera and D. Lubda (Darmstadt, Germany) 433

Separation of ferrocenylethanol enantiomers by microcolumn liquid chromatography with β-cyclodextrin as mobile phase additive
by T. Takeuchi and T. Miwa (Gifu, Japan) 439

Direct enantiomeric resolution of monoterpene hydrocarbons via reversed-phase high-performance liquid chromatography with an α-cyclodextrin bonded stationary phase
by D.W. Armstrong and J. Zukowski (Rolla, MO, USA) 445

One-pot preparation method for a uniform-sized polymer-based chiral stationary phase for high-performance liquid chromatography with polymethacrylamide as a chiral selector
by K. Hosoya, K. Yoshizako, N. Tanaka, K. Kimata and T. Araki (Kyoto, Japan) and J.M.J. Fréchet (Ithaca, NY, USA) 449

Enantiomer separation by high-performance liquid chromatography with (*R*,*R*)-tartaric acid mono-amide derivatives as bifunctional chiral selectors
by N. Ôi, H. Kitahara and F. Aoki (Osaka, Japan) 457

Effect of sample derivatization in liquid chromatographic separations of amine and amino acid enantiomers using diamide-type stationary phases
by K. Sato, H. Nakano and T. Hobo (Tokyo, Japan) 463

Direct resolution of optically active isomers on chiral packings containing ergoline skeletons. II. Enantioseparation of carboxylic acids
by M. Sinibaldi (Rome, Italy), M. Flieger (Prague, Czech Republic), L. Cvak (Opava, Czech Republic) and A. Messina and A. Pichini (Rome, Italy) 471

Analytical and preparative separation of the enantiomers of pyrazole phenyl ether herbicides on three chiral stationary phases
by B.C. Hamper, D.R. Dukesherer and K. Moedritzer (St. Louis, MO, USA) 479

Sensitive enantiomeric separation of aliphatic and aromatic amines using aromatic anhydrides as non-chiral derivatizing agents
by M. Pawlowska, J. Zukowski and D.W. Armstrong (Rolla, MO, USA) 485

Theoretical considerations for chiral recognition

Theoretical studies of brush-type chiral stationary phases (Review)
by K.B. Lipkowitz (Indianapolis, IN, USA) 493

Chromatographic separation of amide diastereomers: correlation with molecular descriptors
by G. Hansson (Göteborg, Sweden) and M. Ahnoff (Mölndal, Sweden) 505

Resolution of non-racemic mixtures in achiral chromatographic systems: a model for the enantioselective effects observed
by E. Gil-Av (Rehovot, Israel) and V. Schurig (Tübingen, Germany) 519

Chiral discrimination of complexes between benzyloxycarbonylglycyl-L-proline and 4-hydroxy-2-dipropylaminoindan in ion-pair chromatography. A molecular mechanics study
by K. Luthman, A.V. Jensen, U. Hacksell, A. Karlsson and C. Pettersson (Uppsala, Sweden) 527

Determination of optical purity by high-performance liquid chromatography upon non-chiral stationary phases with dual circular dichroism/absorption detection
by C. Bertucci, P. Salvadori and L.F. Lopes Guimaraes (Pisa, Italy) 535

Stereochemical features of the stereoselective interaction between (*R*)- or (*S*)-7,7′-bis(1-propen-3-oxy)-2,2′-dihydroxy-1,1′-binaphthyl and quinine
by G. Uccello-Barretta, D. Pini, C. Rosini and P. Salvadori (Pisa, Italy) 541

Mass spectral and semi-empirical studies on chiral discrimination in gaseous aggregation products of protonated dialkyl tartrates
by F.J. Winkler and R. Medina (Freising, Germany) and J. Winkler and H. Krause (Garching, Germany) 549

CHIRBASE, a molecular database for the separation of enantiomers by chromatography
by B. Koppenhoefer, R. Graf, H. Holzschuh, A. Nothdurft and U. Trettin (Tübingen, Germany) and P. Piras and C. Roussel (Marseille, France) 557

Special aspects of separating enantiomers on preparative scale

Contribution of preparative chromatographic resolution to the investigation of chiral phenomena (Review)
by E. Francotte (Basle, Switzerland) 565

Displacement chromatography on cyclodextrin silicas. V. Separation of the enantiomers of 5,10-dideazatetrahydrofolic acid
by L.H. Irgens, G. Farkas and G. Vigh (College Station, TX, USA) 603

Preparative enantiomer separation with modified cyclodextrins as chiral stationary phases
by I. Hardt and W.A. König (Hamburg, Germany) 611

Preparative enantiomer separation of enflurane and isoflurane by inclusion gas chromatography
by V. Schurig and H. Grosenick (Tübingen, Germany) 617

Preparative chromatographic resolution of racemates on chiral stationary phases on laboratory and production scales by closed-loop recycling chromatography
by J. Dingenen (Beerse, Belgium) and J.N. Kinkel (Darmstadt, Germany) 627

AUTHOR INDEX 651